

References

phi :

2010 (2)

Jansen FH, van Schaik RH, Kurstjens J, Horninger W, Klocker H, Bektic J, et al. Prostate-specific antigen (PSA) isoform p2PSA in combination with total PSA and free PSA improves diagnostic accuracy in prostate cancer detection. *European urology*. 2010;57(6):921-7.

Semjonow A, Kopke T, Eltze E, Pepping-Schefers B, Burgel H, Darte C. Pre-analytical in-vitro stability of [-2]proPSA in blood and serum. *Clin Biochem*. 2010;43(10-11):926-8.

2011 (1)

Isharwal S, Makarov DV, Sokoll LJ, Landis P, Marlow C, Epstein JI, et al. ProPSA and diagnostic biopsy tissue DNA content combination improves accuracy to predict need for prostate cancer treatment among men enrolled in an active surveillance program. *Urology*. 2011;77(3):763 e1-6.

2012 (4)

Guazzoni G, Lazzeri M, Nava L, Lughezzani G, Larcher A, Scattoni V, et al. Preoperative prostate-specific antigen isoform p2PSA and its derivatives, %p2PSA and prostate health index, predict pathologic outcomes in patients undergoing radical prostatectomy for prostate cancer. *European urology*. 2012;61(3):455-66.

Lazzeri M, Briganti A, Scattoni V, Lughezzani G, Larcher A, Gadda GM, et al. Serum index test %[-2]proPSA and Prostate Health Index are more accurate than prostate specific antigen and %fPSA in predicting a positive repeat prostate biopsy. *The Journal of urology*. 2012;188(4):1137-43.

Sokoll LJ, Chan DW, Klee GG, Roberts WL, van Schaik RH, Arockiasamy DA, et al. Multi-center analytical performance evaluation of the Access Hybritech(R) p2PSA immunoassay. *Clin Chim Acta*. 2012;413(15-16):1279-83.

Tosoian JJ, Loeb S, Feng Z, Isharwal S, Landis P, Elliot DJ, et al. Association of [-2]proPSA with biopsy reclassification during active surveillance for prostate cancer. *The Journal of urology*. 2012;188(4):1131-6.

2013 (2)

Lamy PJ, Montels F, Tosi D, Leizour B, Bascoul-Mollevis C, Castan F, et al. [Evaluation of (-2)proPSA in combination with total PSA and free PSA for the early detection of prostate cancer]. *Ann Biol Clin (Paris)*. 2013;71(5):537-44.

Scattoni V, Lazzeri M, Lughezzani G, De Luca S, Passera R, Bollito E, et al. Head-to-head comparison of prostate health index and urinary PCA3 for predicting cancer at initial or repeat biopsy. *The Journal of urology*. 2013;190(2):496-501.

2014 (9)

Eminaga O, Bogemann M, Breil B, Titze U, Wotzel F, Eltze E, et al. Preoperative prostate-specific antigen isoform p2PSA \leq 22.5 pg/ml predicts advanced prostate cancer in patients undergoing radical prostatectomy. *Urologic oncology*. 2014;32(8):1317-26.

- Hirama H, Sugimoto M, Ito K, Shiraishi T, Kakehi Y. The impact of baseline [-2]proPSA-related indices on the prediction of pathological reclassification at 1 year during active surveillance for low-risk prostate cancer: the Japanese multicenter study cohort. *J Cancer Res Clin Oncol*. 2014;140(2):257-63.
- Igawa T, Takehara K, Onita T, Ito K, Sakai H. Stability of [-2]Pro-PSA in whole blood and serum: analysis for optimal measurement conditions. *J Clin Lab Anal*. 2014;28(4):315-9.
- Lazzeri M, Abrate A, Lughezzani G, Gadda GM, Freschi M, Mistretta F, et al. Relationship of chronic histologic prostatic inflammation in biopsy specimens with serum isoform [-2]proPSA (p2PSA), %p2PSA, and prostate health index in men with a total prostate-specific antigen of 4-10 ng/ml and normal digital rectal examination. *Urology*. 2014;83(3):606-12.
- Lughezzani G, Lazzeri M, Haese A, McNicholas T, de la Taille A, Buffi NM, et al. Multicenter European external validation of a prostate health index-based nomogram for predicting prostate cancer at extended biopsy. *European urology*. 2014;66(5):906-12.
- Mearini L, Ferri C, Lazzeri M, Bini V, Nunzi E, Fiorini D, et al. Evaluation of prostate-specific antigen isoform p2PSA and its derivatives, %p2PSA, prostate health index and prostate dimension-adjusted related index in the detection of prostate cancer at first biopsy: an exploratory, prospective study. *Urol Int*. 2014;93(2):135-45.
- Na R, Ye D, Liu F, Chen H, Qi J, Wu Y, et al. Performance of serum prostate-specific antigen isoform [-2]proPSA (p2PSA) and the prostate health index (phi) in a Chinese hospital-based biopsy population. *Prostate*. 2014;74(15):1569-75.
- Tallon L, Luangphakdy D, Ruffion A, Colombel M, Devonec M, Champetier D, et al. Comparative evaluation of urinary PCA3 and TMPRSS2: ERG scores and serum phi in predicting prostate cancer aggressiveness. *Int J Mol Sci*. 2014;15(8):13299-316.
- Wang W, Wang M, Wang L, Adams TS, Tian Y, Xu J. Diagnostic ability of %p2PSA and prostate health index for aggressive prostate cancer: a meta-analysis. *Sci Rep*. 2014;4:5012.
- 2015 (8)
- Cantiello F, Russo GI, Ferro M, Cicione A, Cimino S, Favilla V, et al. Prognostic accuracy of Prostate Health Index and urinary Prostate Cancer Antigen 3 in predicting pathologic features after radical prostatectomy. *Urologic oncology*. 2015;33(4):163 e15-23.
- de la Calle C, Patil D, Wei JT, Scherr DS, Sokoll L, Chan DW, et al. Multicenter Evaluation of the Prostate Health Index to Detect Aggressive Prostate Cancer in Biopsy Naive Men. *The Journal of urology*. 2015;194(1):65-72.
- Ferro M, Lucarelli G, Bruzzese D, Perdona S, Mazzarella C, Perruolo G, et al. Improving the prediction of pathologic outcomes in patients undergoing radical prostatectomy: the value of prostate cancer antigen 3 (PCA3), prostate health index (phi) and sarcosine. *Anticancer Res*. 2015;35(2):1017-23.
- Foley RW, Gorman L, Sharifi N, Murphy K, Moore H, Tuzova AV, et al. Improving multivariable prostate cancer risk assessment using the Prostate Health Index. *BJU international*. 2015.
- Fossati N, Buffi NM, Haese A, Stephan C, Larcher A, McNicholas T, et al. Preoperative Prostate-specific Antigen Isoform p2PSA and Its Derivatives, %p2PSA and Prostate Health Index, Predict Pathologic Outcomes in Patients Undergoing Radical Prostatectomy for Prostate Cancer: Results from a Multicentric European Prospective Study. *European urology*. 2015;68(1):132-8.

Loeb S, Sanda MG, Broyles DL, Shin SS, Bangma CH, Wei JT, et al. The prostate health index selectively identifies clinically significant prostate cancer. *The Journal of urology*. 2015;193(4):1163-9.

Mearini L, Nunzi E, Ferri C, Bellezza G, Lolli C, Porrozzi C, et al. Use of the Prostate Health Index for the Detection of Aggressive Prostate Cancer at Radical Prostatectomy. *Urol Int*. 2015;95(4):390-9.

Nordstrom T, Vickers A, Assel M, Lilja H, Gronberg H, Eklund M. Comparison Between the Four-kallikrein Panel and Prostate Health Index for Predicting Prostate Cancer. *European urology*. 2015;68(1):139-46.

2016 (1)

Boegemann M, Stephan C, Cammann H, Vincendeau S, Houlgatte A, Jung K, et al. The percentage of prostate-specific antigen (PSA) isoform [-2]proPSA and the Prostate Health Index improve the diagnostic accuracy for clinically relevant prostate cancer at initial and repeat biopsy compared with total PSA and percentage free PSA in men aged ≤ 65 years. *BJU international*. 2016;117(1):72-9.

4Kscore : références bibliographiques

2008 (1)

Vickers AJ, Cronin AM, Aus G, Pihl CG, Becker C, Pettersson K, et al. A panel of kallikrein markers can reduce unnecessary biopsy for prostate cancer: data from the European Randomized Study of Prostate Cancer Screening in Goteborg, Sweden. *BMC Med*. 2008;6:19.

2010 (5)

Benchikh A, Savage C, Cronin A, Salama G, Villers A, Lilja H, et al. A panel of kallikrein markers can predict outcome of prostate biopsy following clinical work-up: an independent validation study from the European Randomized Study of Prostate Cancer screening, France. *BMC Cancer*. 2010;10:635.

Gupta A, Roobol MJ, Savage CJ, Peltola M, Pettersson K, Scardino PT, et al. A four-kallikrein panel for the prediction of repeat prostate biopsy: data from the European Randomized Study of Prostate Cancer screening in Rotterdam, Netherlands. *Br J Cancer*. 2010;103(5):708-14.

Vickers A, Cronin A, Roobol M, Savage C, Peltola M, Pettersson K, et al. Reducing unnecessary biopsy during prostate cancer screening using a four-kallikrein panel: an independent replication. *J Clin Oncol*. 2010;28(15):2493-8.

Vickers AJ, Cronin AM, Aus G, Pihl CG, Becker C, Pettersson K, et al. Impact of recent screening on predicting the outcome of prostate cancer biopsy in men with elevated prostate-specific antigen: data from the European Randomized Study of Prostate Cancer Screening in Gothenburg, Sweden. *Cancer*. 2010;116(11):2612-20.

Vickers AJ, Cronin AM, Roobol MJ, Savage CJ, Peltola M, Pettersson K, et al. A four-kallikrein panel predicts prostate cancer in men with recent screening: data from the European Randomized Study of Screening for Prostate Cancer, Rotterdam. *Clin Cancer Res*. 2010;16(12):3232-9.

2013 (1)

Carlsson S, Maschino A, Schroder F, Bangma C, Steyerberg EW, van der Kwast T, et al. Predictive value of four kallikrein markers for pathologically insignificant compared with aggressive prostate cancer in radical prostatectomy specimens: results from the European Randomized

Study of Screening for Prostate Cancer section Rotterdam. *European urology*. 2013;64(5):693-9.

2015 (5)

Braun K, Sjoberg DD, Vickers AJ, Lilja H, Bjartell AS. A Four-kallikrein Panel Predicts High-grade Cancer on Biopsy: Independent Validation in a Community Cohort. *European urology*. 2015.

Bryant RJ, Sjoberg DD, Vickers AJ, Robinson MC, Kumar R, Marsden L, et al. Predicting high-grade cancer at ten-core prostate biopsy using four kallikrein markers measured in blood in the ProtecT study. *J Natl Cancer Inst*. 2015;107(7).

Nordstrom T, Vickers A, Assel M, Lilja H, Gronberg H, Eklund M. Comparison Between the Four-kallikrein Panel and Prostate Health Index for Predicting Prostate Cancer. *European urology*. 2015;68(1):139-46.

Parekh DJ, Punnen S, Sjoberg DD, Asroff SW, Bailen JL, Cochran JS, et al. A multi-institutional prospective trial in the USA confirms that the 4Kscore accurately identifies men with high-grade prostate cancer. *European urology*. 2015;68(3):464-70.

Stattin P, Vickers AJ, Sjoberg DD, Johansson R, Granfors T, Johansson M, et al. Improving the Specificity of Screening for Lethal Prostate Cancer Using Prostate-specific Antigen and a Panel of Kallikrein Markers: A Nested Case-Control Study. *European urology*. 2015;68(2):207-13.

MiPS : références bibliographiques

2011 (1)

Tomlins SA, Aubin SM, Siddiqui J, Lonigro RJ, Sefton-Miller L, Miick S, et al. Urine TMPRSS2:ERG fusion transcript stratifies prostate cancer risk in men with elevated serum PSA. *Sci Transl Med*. 2011;3(94):94ra72.

2013 (2)

Cornu JN, Cancel-Tassin G, Egrot C, Gaffory C, Haab F, Cussenot O. Urine TMPRSS2:ERG fusion transcript integrated with PCA3 score, genotyping, and biological features are correlated to the results of prostatic biopsies in men at risk of prostate cancer. *Prostate*. 2013;73(3):242-9.

Lin DW, Newcomb LF, Brown EC, Brooks JD, Carroll PR, Feng Z, et al. Urinary TMPRSS2:ERG and PCA3 in an active surveillance cohort: results from a baseline analysis in the Canary Prostate Active Surveillance Study. *Clin Cancer Res*. 2013;19(9):2442-50.

2014 (2)

Leyten GH, Hessels D, Jannink SA, Smit FP, de Jong H, Cornel EB, et al. Prospective multicentre evaluation of PCA3 and TMPRSS2-ERG gene fusions as diagnostic and prognostic urinary biomarkers for prostate cancer. *European urology*. 2014;65(3):534-42.

Tallon L, Luangphakdy D, Ruffion A, Colombel M, Devonec M, Champetier D, et al. Comparative evaluation of urinary PCA3 and TMPRSS2: ERG scores and serum phi in predicting prostate cancer aggressiveness. *Int J Mol Sci*. 2014;15(8):13299-316.

2015 (1)

Tomlins SA, Day JR, Lonigro RJ, Hovelson DH, Siddiqui J, Kunju LP, et al. Urine TMPRSS2:ERG Plus PCA3 for Individualized Prostate Cancer Risk Assessment. *European urology*. 2015.

GPS : références bibliographiques

2013 (1)

Knezevic D, Goddard AD, Natraj N, Cherbavaz DB, Clark-Langone KM, Snable J, et al. Analytical validation of the Oncotype DX prostate cancer assay - a clinical RT-PCR assay optimized for prostate needle biopsies. *BMC Genomics*. 2013;14:690.

2014 (1)

Klein EA, Cooperberg MR, Magi-Galluzzi C, Simko JP, Falzarano SM, Maddala T, et al. A 17-gene assay to predict prostate cancer aggressiveness in the context of Gleason grade heterogeneity, tumor multifocality, and biopsy undersampling. *European urology*. 2014;66(3):550-60.

2015 (1)

Cullen J, Rosner IL, Brand TC, Zhang N, Tsiatis AC, Moncur J, et al. A Biopsy-based 17-gene Genomic Prostate Score Predicts Recurrence After Radical Prostatectomy and Adverse Surgical Pathology in a Racially Diverse Population of Men with Clinically Low- and Intermediate-risk Prostate Cancer. *European urology*. 2015;68(1):123-31.

Prolaris : références bibliographiques

2011 (1)

Cuzick J, Swanson GP, Fisher G, Brothman AR, Berney DM, Reid JE, et al. Prognostic value of an RNA expression signature derived from cell cycle proliferation genes in patients with prostate cancer: a retrospective study. *Lancet Oncol*. 2011;12(3):245-55.

2012 (1)

Cuzick J, Berney DM, Fisher G, Mesher D, Moller H, Reid JE, et al. Prognostic value of a cell cycle progression signature for prostate cancer death in a conservatively managed needle biopsy cohort. *Br J Cancer*. 2012;106(6):1095-9.

2013 (2)

Cooperberg MR, Simko JP, Cowan JE, Reid JE, Djilivand A, Bhatnagar S, et al. Validation of a cell-cycle progression gene panel to improve risk stratification in a contemporary prostatectomy cohort. *J Clin Oncol*. 2013;31(11):1428-34.

Freedland SJ, Gerber L, Reid J, Welbourn W, Tikishvili E, Park J, et al. Prognostic utility of cell cycle progression score in men with prostate cancer after primary external beam radiation therapy. *Int J Radiat Oncol Biol Phys*. 2013;86(5):848-53.

2014 (2)

Arsov C, Jankowiak F, Hiester A, Rabenalt R, Quentin M, Schimmoller L, et al. Prognostic value of a cell-cycle progression score in men with prostate cancer managed with active surveillance after MRI-guided prostate biopsy--a pilot study. *Anticancer Res*. 2014;34(5):2459-66.

Bishoff JT, Freedland SJ, Gerber L, Tennstedt P, Reid J, Welbourn W, et al. Prognostic utility of the cell cycle progression score generated from biopsy in men treated with prostatectomy. *The Journal of urology*. 2014;192(2):409-14.

2015 (1)

Cuzick J, Stone S, Fisher G, Yang ZH, North BV, Berney DM, et al. Validation of an RNA cell cycle progression score for predicting death from prostate cancer in a conservatively managed needle biopsy cohort. *Br J Cancer*. 2015;113(3):382-9.

Decipher : références bibliographiques

2013 (2)

Erho N, Crisan A, Vergara IA, Mitra AP, Ghadessi M, Buerki C, et al. Discovery and validation of a prostate cancer genomic classifier that predicts early metastasis following radical prostatectomy. *PLoS One*. 2013;8(6):e66855.

Karnes RJ, Bergstralh EJ, Davicioni E, Ghadessi M, Buerki C, Mitra AP, et al. Validation of a genomic classifier that predicts metastasis following radical prostatectomy in an at risk patient population. *The Journal of urology*. 2013;190(6):2047-53.

2014 (2)

Den RB, Feng FY, Showalter TN, Mishra MV, Trabulsi EJ, Lallas CD, et al. Genomic prostate cancer classifier predicts biochemical failure and metastases in patients after postoperative radiation therapy. *Int J Radiat Oncol Biol Phys*. 2014;89(5):1038-46.

Ross AE, Feng FY, Ghadessi M, Erho N, Crisan A, Buerki C, et al. A genomic classifier predicting metastatic disease progression in men with biochemical recurrence after prostatectomy. *Prostate Cancer Prostatic Dis*. 2014;17(1):64-9.

2015 (3)

Cooperberg MR, Davicioni E, Crisan A, Jenkins RB, Ghadessi M, Karnes RJ. Combined value of validated clinical and genomic risk stratification tools for predicting prostate cancer mortality in a high-risk prostatectomy cohort. *European urology*. 2015;67(2):326-33.

Den RB, Yousefi K, Trabulsi EJ, Abdollah F, Choerung V, Feng FY, et al. Genomic classifier identifies men with adverse pathology after radical prostatectomy who benefit from adjuvant radiation therapy. *J Clin Oncol*. 2015;33(8):944-51.

Klein EA, Yousefi K, Haddad Z, Choerung V, Buerki C, Stephenson AJ, et al. A genomic classifier improves prediction of metastatic disease within 5 years after surgery in node-negative high-risk prostate cancer patients managed by radical prostatectomy without adjuvant therapy. *European urology*. 2015;67(4):778-86.

2016 (1)

Ross AE, Johnson MH, Yousefi K, Davicioni E, Netto GJ, Marchionni L, et al. Tissue-based Genomics Augments Post-prostatectomy Risk Stratification in a Natural History Cohort of Intermediate- and High-Risk Men. *European urology*. 2016;69(1):157-65.